

Pressure driven neutral-ionic transition in ClMePD-DMeDCNQI

Luca Farina and Aldo Brillante

Dip. di Chimica Fisica ed Inorganica, Università di Bologna, Viale Risorgimento 4, 40136-I Bologna, Italy

Matteo Masino and Alberto Girlando

Dip. Chimica Generale ed Inorganica, Chimica Analitica e Chimica Fisica, Università di Parma, Parco Area delle Scienze, 43100-I Parma, Italy

Application of about 0.8 GPa pressure is sufficient to induce the neutral-ionic transition in the mixed stack charge-transfer crystal 2-chloro-5-methyl-*p*-phenylenediamine-2,5-dimethyldicyanoquinonediimine (ClMePD-DMeDCNQI). The ionicity increases continuously from ~ 0.35 at ambient conditions to ~ 0.65 when the pressure is raised up to 2 GPa. Moreover, stack dimerization begins well before the crossing of the neutral-ionic interface. The evolution of the transition is similar to what observed in the temperature induced phase change in the same compound. A distinguishing feature is represented by the simultaneous presence of domains of molecules with slightly different ionicities across the transition pressure. A comparison of the present example of pressure driven neutral-ionic transition with the well studied cases of tetrathiafulvalene-chloranil and of tetrathiafulvalene-2,5-dichloro-*p*-benzoquinone puts in evidence the remarkably different evolution of the three transitions.

I. INTRODUCTION

Neutral-ionic (NI) phase transitions in mixed stack charge transfer (CT) crystals¹ have recently received renewed interest due to the expectation of unusual physical properties at the NI borderline. For instance, exact diagonalization studies have suggested the possible existence of a metallic state just at the point marking the passage from N to I ground state of a regular mixed stack chain.²⁻⁴ Metallic behavior around this particular point has been indeed discovered.⁵ Moreover, a large increase of the dielectric constant, associated with the collective transfer of charge, has been found in proximity of the NI transition.⁶ Finally, systems on the I side show strong tendency towards dimerization, due to the Peierls instability,³ and may therefore display a ferroelectric ground state.⁷ To explore such phenomena, it is useful to investigate CT crystals that undergo a continuous ionicity change across the NI borderline. Several years ago our group reported an example of continuous ionicity change under pressure for tetrathiafulvalene-2,5-dichloro-*p*-benzoquinone (TTF-DCIBQ).⁸ More recently, a first example of a continuous NI transition induced by lowering temperature has been observed for 2-chloro-5-methyl-*p*-phenylenediamine-2,5-dimethyldicyanoquinonediimine (ClMePD-DMeDCNQI).⁹ A detailed characterization of this transition has shown that there is a sort of feedback mechanism between the Peierls dimerization transition and the increase of ionicity at the NI interface.¹⁰

In the present paper we investigate the NI phase transition in ClMePD-DMeDCNQI induced by pressure at room temperature. We find that the pressure induced phase transition in ClMePD-DMeDCNQI is somewhat different from the transition induced by temperature.¹⁰ Furthermore, in ClMePD-DMeDCNQI

the ionicity change across the NI borderline is practically continuous, as in TTF-DCIBQ,⁸ but the evolution of the two transitions is quite different. We also make comparison with the NI phase transition of the prototype compound tetrathiafulvalene-chloranil (TTF-CA), which is known to exhibit a discontinuous NI phase transition both under pressure^{11,12} and under temperature.^{13,14}

II. EXPERIMENTAL

ClMePD-DMeDCNQI crystals have been prepared by mixing saturated solutions in dichloromethane followed by slow evaporation of the solvent.⁹ Infrared (IR) spectra have been recorded with a Nicolet Nexus 470 FTIR on powdered samples loaded in a diamond-anvil cell (DAC) with nujol or perfluorocarbons (PFC) as pressure media. High pressure Raman spectra were obtained with a Renishaw System 1000 Microscope, with excitation from an Ar ion laser ($\lambda = 514.5$ nm). A tiny crystal was loaded in the DAC using nujol as pressure medium. The spectral resolution in both IR and Raman spectra is 2 cm^{-1} . Pressure calibration was done with the ruby luminescence technique.¹⁵ Accuracy of the pressure reading was about 0.05 GPa in the pressure range up to 3 GPa.

III. RESULTS

ClMePD-DMeDCNQI crystallizes in the triclinic system, space group P1, with $a=7.463\text{\AA}$, $b=7.504\text{\AA}$, $c=7.191\text{\AA}$, $\alpha=91.23^\circ$, $\beta=112.19^\circ$, $\gamma=96.91^\circ$, and $Z=1$.⁹ The electron donor (D) ClMePD and the electron acceptor (A) DMeDCNQI form mixed stack columns along the *b* axis. The crystal does not have symmetry elements, and the molecules are located in general positions. Therefore the stack structure is not constrained by symmetry.




FIG. 1. Pressure dependence of the IR and Raman spectra of ClMePD-DMeDCNQI in the 1525-1700 cm^{-1} spectral region. The pressures are indicated on the right side of the figure.

Within the accuracy of the X-ray analysis,⁹ at room temperature the stack is regular, that is, each molecule has equal distance (CT integral) with its two nearest neighbors along the chain. However, the stack structure presents disorder in the relative orientation of the two molecules: the 2- and 5- substitutional sites of ClMePD are considered to be occupied by chlorine and methyl group with equal probability along the stack.

As it is well known,¹³ vibrational spectroscopy is a very useful tool to study NI phase transitions of mixed stack CT crystals. Vibrational frequencies respond to molecular charge variations, therefore yielding a method to directly estimate the degree of ionicity ρ . Moreover, the interaction between the CT electrons and the molecular vibrations (e - mv coupling) allows one to discriminate regular from dimerized stack structure: The stack dimerization is marked by the appearance in the IR spectra of the (Raman active) totally symmetric modes, which are forbidden in the case of regular stack.^{16,17} The IR intensity of these modes is related to the extent of dimerization, and can be taken as the order parameter of the dimerization phase transition.




FIG. 2. Pressure dependence of the IR and Raman spectra of ClMePD-DMeDCNQI in the 800-1300 cm^{-1} spectral region. The pressures are indicated on the right side of the figure.

In this paper we only deal with the spectral features directly connected with the phase transition, as singled out in ref. 10, where a satisfactory and rather complete interpretation of the IR and Raman spectra has been achieved. Figs. 1 and 2 report the IR and Raman spectra as function of pressure, in the spectral regions 1525-1700 and 800-1300 cm^{-1} , respectively. The IR active mode DMeDCNQI $b_u\nu_{46}$ mode (1570 cm^{-1} at ambient conditions, marked A in the fig. 1) has been chosen as the primary probe of ionicity, since it shows a large ionization frequency shift (61 cm^{-1}).¹⁰ The other DMeDCNQI mode showing a linear dependence on ρ , the $b_u\nu_{47}$, cannot be used as a secondary check of the ionicity,¹⁰ since in the lack of polarized IR data we cannot try to disentangle this mode in the very complex spectral region 1480-1520 cm^{-1} .

By increasing the pressure the DMeDCNQI $b_u\nu_{46}$ (A band) shifts towards lower frequencies from 1570 to 1550 cm^{-1} , indicating an increase in the molecular ionicity. We notice, however, that already at 0.22 GPa a dip appears on the low-frequency side of the A band. This spectral feature (marked A' in fig. 1) gradually gains

intensity as the pressure increases. Between 0.76 and 0.99 GPa its intensity gets comparable to that of the A band, and when pressure is raised up to 1.57 GPa the A band merges into the A' band. Since only the DMeDCNQI $b_u\nu_{46}$ mode is expected in this frequency region, we interpret the A and A' bands as due to this mode, in correspondence with species of different ionicities. Thus the simultaneous presence of A and A' bands signals the presence of domains of slightly different ionicities in the crystal. This kind of behavior, which would suggest staging in the phase transition, is not observed in the temperature induced transition, where coexistence of phases at different ionicities can safely be excluded.^{9,10} However, the behavior reported in fig. 1 varies on changing sample during different pressure runs, showing slightly different profiles with respect to those reported in fig. 1. In any case the structure of the A band is always rather complex showing in some cases appreciable intensity of the A' band already at very low pressures. The problem might be accounted for the different samples used in IR spectra, i.e. powdered crystals dispersed in nujol or PFC mulls. As already observed in the NI transition of TTF-CA,^{12,13} defects and imperfections may indeed affect the course of the phase transition, and measurements on single crystals are needed to safely ascertain the presence of staging. In fig. 3, upper panel, we report the pressure dependence of the $b_u\nu_{46}$ frequency, taken as the peak frequency of the most intense among the A, A' bands.

From the pressure evolution of IR spectra in figs. 1 and 2, one easily recognizes the presence of $e-mv$ induced bands, which signal the onset of stack dimerization. We focus on the two $e-mv$ induced IR bands at 1587 and 1241 cm^{-1} , assigned to DMeDCNQI totally-symmetric modes (marked by an asterisk in figs. 1 and 2). The pressure dependence of their intensity, divided by the intensity of the normally IR active DMeDCNQI $b_u\nu_{51}$ band (at 1285 cm^{-1}), are reported in the lower panel of fig. 3. As previously reported,^{9,10} at ambient conditions the stack is already dimerized to some extent, as shown by the presence of $e-mv$ induced bands in the ambient pressure IR spectrum. By increasing the pressure, the intensity of these bands increases gradually, indicating an increase in the extent of dimerization. We remark that by increasing the pressure also several Raman bands assigned to totally-symmetric modes of DMeDCNQI gain appreciable intensity over the bands of ClMePD. This unexpected behavior is clearly evident by looking at the pressure evolution of the already mentioned DMeDCNQI $a_g\nu_{10}$ and ν_5 modes in fig.1. A possible explanation can be formulated on the basis of the ionicity variation under pressure. In fact, by increasing the degree of ionicity of ClMePD-DMeDCNQI, either by lowering temperature or by increasing pressure, a band due to the lowest intramolecular electronic excitation of DMeDCNQI^- appears around 2.4 eV in the visible absorption spectra.⁹ Since the Raman exciting line used in the present work is fully in resonance with this band ($\lambda = 514.5 \text{ nm} = 2.41 \text{ eV}$),




FIG. 3. Upper panel: Pressure dependence of the frequency of the DMeDCNQI $b_u\nu_{46}$ mode. Diamonds and circles refer to nujol and perfluorocarbon spectra, respectively. Lower panel: Pressure dependence of the relative intensities of the $e-mv$ induced bands corresponding to the DMeDCNQI $a_g\nu_5$ (triangles) and ν_{10} modes (inverted triangles).

we observe resonance intensity enhancement of the Raman bands due to DMeDCNQI molecular units.

Before discussing in detail the pressure induced NI phase transition in ClMePD-DMeDCNQI one last observation on vibrational spectra is in order. On the basis of the vibronic model^{16,17} for dimerized mixed stack CT crystals, $e-mv$ induced IR bands and their Raman counterparts should display frequency coincidence. On the contrary, we observe appreciable frequency differences in the high pressure spectra, the most noticeable one being observed for the DMeDCNQI $a_g\nu_5$ band, which is present at 1585 cm^{-1} in the Raman spectra at 1.53 GPa and at 1598 cm^{-1} in the IR spectra at 1.57 GPa (fig. 1). The

origin of this effect might be again attributed to the above mentioned resonance Raman conditions. Resonance effects on Raman spectra under pressure can indeed single out domains of different ionicities within the crystal, so that the resulting peak maxima may differ from those of the IR spectra. Furthermore, one has to keep in mind that different experimental conditions have been used in measuring Raman and IR spectra, namely single crystal for Raman, and powdered samples for IR.

IV. DISCUSSION

We first discuss the ionicity change of ClMePD-DMeDCNQI crystal under pressure. Assuming that the frequency of the DMeDCNQI $b_u\nu_{46}$ varies linearly with the degree of ionicity,¹⁸ from the upper panel of fig. 3 we immediately derive the pressure dependence of ρ reported in the left panel of fig. 4. The ionicity increases smoothly with pressure from ~ 0.35 at ambient conditions to ~ 0.65 at 3 GPa. In fig. 4, we have limited the pressure to 2.2 GPa, as the frequency of the $b_u\nu_{46}$ band saturates above such pressure (fig. 3, upper panel). We remark that to estimate ρ we have used the frequency of the most intense among the A,A' bands of fig. 1, so that fig. 4 reports the ionicity of the prevailing phase, and the possible staging of the transition is not evidenced. The difference in frequency between A and A' bands is of the order of 4 cm^{-1} , corresponding to a ρ difference of the order of 0.07. The overall ionicity variation is about 0.3 in the pressure range from ambient to 2 GPa. We note that this value should be considered a lower limit, as we have not corrected for the general frequency increase due

to compression effects. We have increased the pressure up to 5.9 GPa, confirming that the ionicity apparently saturates around 2 GPa. However, above 3 GPa the uncertainty in the estimate of ρ increases, as the frequency hardening due to anharmonicity cannot be disregarded anymore.⁸

In the left panel of fig. 4 we also report the pressure dependence of the intensity of the $e-mv$ induced IR bands corresponding to the DMeDCNQI $a_g\nu_5$ and ν_{10} modes (fig. 3, lower panel). For convenience, we have normalized the intensities of both bands to the value at 3 GPa. From the figure it is seen that the intensity of these bands, already present at ambient conditions, gradually increases by increasing pressure, reaching a plateau at about 1.3 GPa. The maximum slope is around 0.5 GPa. As in the temperature induced phase transition,¹⁰ the dimerization occurs well before the crossing of the NI borderline. On the other hand, the already mentioned simultaneous presence of domains of different ionicities (staging in the phase transition) suggests that the pressure induced phase transition occurs with a mechanism different from that of the temperature induced phase transition, which is a truly continuous and second order transition. The peculiarity of the NI transition in ClMePD-DMeDCNQI is that the coupling between a first order phase transition (order parameter: the degree of ionicity) and a second order, displacive one (order parameter: the degree of dimerization) is particularly evident.¹⁰

The above interplay between first and second order phase transition is well appreciated if we compare the pressure induced NI transition in ClMePD-DMeDCNQI




FIG. 4. Pressure dependence of the ionicity (ρ) and of the normalized intensity of the $e-mv$ induced bands of ClMePD-DMeDCNQI (left panel), TTF-DCIBQ (center panel, from ref. 8) and TTF-CA (right panel, adapted from refs. 11,12). The dashed vertical line is a visual aid to approximately separate N and I states (ρ between 0.5 and 0.6).

with the analogous transitions observed in TTF-DCIBQ and in TTF-CA. Fig. 4, center and right panel, reports the corresponding pressure variation of the ionicity, and the normalized intensity of the $e-mv$ induced bands.^{8,11,12} In TTF-DCIBQ we have a continuous ionicity change, as in ClMePD-DMeDCNQI, but the onset of dimerization occurs when ρ is above 0.5, and the NI borderline has been crossed. In TTF-CA, on the other hand, the NI transition is first order, with a ionicity jump of about 0.2 at 1.1 GPa, and again the stack dimerization sets up once the NI borderline is crossed. In both cases, therefore, the transition appears to be driven by the degree of ionicity, followed by the dimerization transition. In the case of ClMePD-DMeDCNQI, on the opposite, the dimerization starts well before the crossover from neutral to ionic regime, causing a strict interplay between dimerization and ionicity. It is worthwhile to remark that in TTF-CA the first order pressure induced phase transition is preceded by a precursor regime (from about 0.7 GPa, not shown in fig. 4), where dimerized quasi-ionic species are present at the same time as the quasi-neutral regular stack.¹² Such pre-transition regime is not present in the temperature induced transition of TTF-CA, at least when well grown single crystals are used.^{13,19} The pre-transition regime has been interpreted as characterized by the presence of thermally fluctuating collective low-energy excitations. However, one cannot exclude that the observed phenomena are instead due to macroscopic, static domains of different ionicities, caused by defects in the powdered samples, as well as by pressure inhomogeneities. The problem is very similar to that encountered in ClMePD-DMeDCNQI, where we observe the coexistence of domains of slightly different ionicities. Such coexistence has not been detected in TTF-DCIBQ.⁸

The quite different behavior of the pressure induced NI phase transitions in ClMePD-DMeDCNQI, TTF-DCIBQ, and TTF-CA, as evidenced in fig. 4, can be rationalized on the basis of existing models of the NI interface. At the basic level, N or I ground states arise from the competition between Δ , the energy difference between donor and acceptor sites, and the energy gain due to the increase of Madelung energy following a lattice contraction. The hopping integral, t , mixes the fully N and I quantum states, and ρ may then change continuously from zero to one. Electron-electron interactions and $e-mv$ coupling, on the other hand, favor first order transitions, with discontinuous jump of ionicity, whereas the modulation of t by the intermolecular, lattice phonons ($e-lph$ coupling) is able to induce the dimerization of the stack, yielding a second order, Peierls-like, phase transition.³

Most of the known NI transitions exhibit a discontinuous jump of ionicity, accompanied by the stack dimerization. For this reason the interplay between ionicity and dimerization has been often overlooked. On the other hand, it has been shown that whereas on the I side the system is intrinsically unstable towards stack dimerization, the dimerization transition may also occur on the




FIG. 5. Asymmetry parameter $\phi \propto (t_i - t_{i+1})/(t_i + t_{i+1})$ as a function of ρ . The calculation is made for the same set of parameters, except for the electron-electron interaction strength, which is zero in the lower panel and different from zero in the upper panel. From ref. 3.

N side, provided the $e-lph$ coupling is strong enough.³ The extent of dimerization and the corresponding energy gain are connected to the ionicity. Fig. 5 shows the calculated ρ dependence of the asymmetry parameter, $\phi \propto (t_i - t_{i+1})/(t_i + t_{i+1})$, directly related to the extent of dimerization.³ The upper panel refers to a set of parameters that yields a continuous ionicity change, whereas in the lower panel the value of the electron-electron interaction gives rise to a jump in ρ (the shaded area represents a forbidden ionicity region). TTF-DCIBQ and TTF-CA then can be roughly described by the upper and lower panel of fig. 5, respectively. In both cases the $e-lph$ coupling strength is not enough to induce the dimerization phase transition before the system is passed on the I side. Thus the transition is driven by the ionicity change, and the dimerization follows. In the case of ClMePD-DMeDCNQI, instead, the $e-lph$ coupling yields the stack dimerization on the N side, when ρ is of the order of 0.35 (fig. 4). Fig. 5 shows that on the N side ϕ increases with ρ , so that the dimerization represents the order parameter of the second order phase transition, but the ρ increase yields in turn to the increase of ϕ . The maximum extent of dimerization, and corre-

spondingly the maximum gain in energy upon dimerization, occurs when ρ is between 0.6 and 0.7, and indeed in ClMePD-DMeDCNQI the pressure dependence of the ionicity shows tendency to saturation (fig. 4).

The above sketched scenario of pressure driven transition in ClMePD-DMeDCNQI is rather simplified, as it does not consider several factors which are currently believed important ingredients of NI phase transitions. Indeed disorder, low-lying collective excitations (CT strings),²⁰ or inter-chain interactions,^{21,22} are not taken into account. Despite of that, we believe our picture catches the salient features of NI phase transition in ClMePD-DMeDCNQI, whereas some aspects still remain to be clarified. From this point of view, the discovery of a new type of NI transition, such as that of ClMePD-DMeDCNQI, where the ionicity change is continuous *and* is preceded by the stack dimerization, represents an additional opportunity for investigating the fascinating multiple facets of pressure and temperature induced NI phase transitions.

ACKNOWLEDGMENTS

We acknowledge helpful discussions with A. Painelli. This work has been supported by the Italian National Research Council (CNR) within its “Progetto Finalizzato Materiali Speciali per tecnologie Avanzate II”, and the Ministry of University and of Scientific and Technological Research (MURST).

- ¹¹ M.Hanfland, A.Brillante, A.Girlando, and K.Syassen, Phys. Rev. B **38**, 1456 (1988).
- ¹² H.Okamoto, T.Koda, Y.Tokura, T.Mitani, and G.Saito Phys. Rev. B **39**, 10693 (1989).
- ¹³ A.Girlando, F.Marzola, C.Pecile, and J.B.Torrance J.Chem.Phys. **79**, 1075 (1983).
- ¹⁴ Y.Tokura, Y.Kaneko, H.Okamoto, S.Tanuma, T.Koda, T.Mitani, and G.Saito, Mol. Cryst. Liq. Cryst., **125**, 71 (1985).
- ¹⁵ G.J.Piermarini, S.Block, J.P.Barnett and R.A.Forman, J.Appl.Phys. **46**, 2774 (1975).
- ¹⁶ C.Pecile, A.Painelli, and A.Girlando Mol. Cryst. Liq. Cryst. **171**, 69 (1989).
- ¹⁷ A.Painelli and A.Girlando, J.Chem.Phys. **84**, 5655 (1986).
- ¹⁸ H.Kobayashi, A.Miyamoto, R.Kato, F.Sakai, A.Kobayashi, Y.Yamakita, Y.Furukawa, M.Tasumi and T.Watanabe, Phys.Rev. B **47**, 3500 (1993).
- ¹⁹ Y.Tokura, Y.Kaneko, H.Okamoto, S.Tanuma, T.Koda, T.Mitani, and G.Saito, Mol. Cryst. Liq. Cryst. **125**, 71 (1985).
- ²⁰ Shin-ya Koshihara, Y.Takahashi, H.Sakai, Y.Tokura, and T.Luty, J.Phys.Chem. B, **103**, 2592 (1999).
- ²¹ E.Collet, M.Buron-LeCointe, M.H.Lemée-Cailleau, H.Cailleau, L.Toupet, M.Meven, S.Mattauch, G.Heger, and N.Karl, Phys.Rev. B, **63**, 054105 (2001).
- ²² V.Oison, C.Katan, and C.Koenig, J. Phys. Chem. A, **105**, 4300 (2001).

-
- ¹ J.B.Torrance, J.E.Vazquez, J.J.Mayerle, and V.Y.Lee, Phys.Rev.Lett. **46**, 253 (1981); J.B.Torrance, A.Girlando, J.J.Mayerle, J.I.Crowley, V.Y.Lee, P.Batail, and S.J.LaPlace, Phys.Rev.Lett. **47**, 1747 (1981).
 - ² A.Painelli and A.Girlando, J.Chem.Phys. **87**, 1705 (1987).
 - ³ A.Painelli and A.Girlando, Phys.Rev. B **37**, 5748 (1988).
 - ⁴ Y.Anusooya-Pati, Z.G.Soos, and A.Painelli, Phys. Rev. B **63**, 205118 (2001).
 - ⁵ G.Saito, Sang-Soo Pac and O.O.Drozдова, Synth. Metals, in press
 - ⁶ S.Horiuchi, R.Kumai, Y.Okimoto, and Y.Tokura, Phys.Rev.Lett. **85**, 5210 (2000).
 - ⁷ M.H.Lemée-Cailleau, M.Le Cointe, H.Cailleau, T.Luty, F.Moussa, J.Roos, D.Brinkmann, B.Toudic, C.Ayache, and N.Karl, Phys.Rev.Lett. **79**, 1690 (1997).
 - ⁸ A. Brillante and A.Girlando, Phys.Rev. B **45**, 7026 (1992); A.Girlando, A.Painelli, C.Pecile, G-L.Calestani, C.Rizzoli and R.M.Metzger, J.Chem.Phys. **98**, 7692 (1993).
 - ⁹ S.Aoki and T.Nakayama, Phys.Rev. B **56**, R2893 (1997).
 - ¹⁰ M.Masino, A.Girlando, L.Farina, and A.Brillante, Phys. Chem. Chem. Phys. **3**, 1904 (2001).